

After the laser

For a few days after the laser you may notice a lot of floaters. The capsule segment drops to the bottom of the eye over a few days, and you may notice large or small floaters at this time; they are normally not troublesome.



New spectacles


You may need a test for new spectacles. Normally this is best carried out at your own optician's office. However, if your sight is bad, ask for an appointment for a test in the eye clinic.

Are there complications from laser?

The laser is considered very safe, with very few complications. Occasionally a little more laser is needed to release the capsule on a second occasion.

Only a small hole in the capsule is needed, so the lens implant still has enough supports

Less often, the laser can disturb the retina, the film lining in the back of the eye. For instance, very rarely a little fluid can build up in the retina.



In addition, the retina may rarely develop a tiny tear months-years later. If this happens you may notice:

- Flashes of light during the daytime
- Floaters in your vision (it is normal to have these in the weeks after the laser)
- A "curtain" drifting across your vision (if the tear extends into a retinal detachment)

If you notice a lot of flashes and floaters all of a sudden, you need to have your eye examined to check if there is a small tear in the retina.

If a "curtain" or "shutter" drifts across your vision, you need to see an ophthalmologist the same day.

As a precaution you should test your sight every day by covering one eye and checking your sight in the other eye.

Dr. _____


Telephone: _____

Next appointment _____

Thickening of the posterior capsule & laser treatment

This leaflet explains:


- *What a thickened capsule is*
- *How your sight is affected*
- *What the laser is like*
- *Other details about laser*



The film behind your lens implant can thicken like frosted glass.

What a cataract is

Normally light enters the eye from the front, passes through a clear lens, and reaches the back of the eye so you can see. If the lens becomes cloudy like frosted glass, light cannot pass through, and you cannot see properly. When the lens becomes cloudy, it is called a 'cataract'.



Cataract surgery


If you have had a cataract operation, a clear plastic lens is placed in the 'bag' left behind when the cataract is removed. The new lens focuses the light onto the back of the eye, just like the original human lens. The 'bag' is called a capsule; the capsule supports the implant.

A thickened posterior capsule

This capsule is normally clear like a glass window. It is very thin, like cellophane. In a small number of patients the capsule thickens up, and becomes a little opaque, like a frosted glass window. This stops the light from reaching the back of the eye.

If this happens, your sight becomes misty, and it can become difficult to see at night, or in sunlight.

Capsule thickening can happen in the months after your cataract operation, but more commonly occurs two or three years after. The capsule thickening does not damage the eye in any way; it merely makes the sight fuzzy.



Laser treatment

Laser treatment makes a small hole in the center of the capsule, and can restore your vision to how it was after the cataract operation. The treatment does not hurt, and you hardly notice anything at all. It takes about fifteen minutes; the laser machine is in the outpatient clinic.

Your physician may or may not use a contact lens for the laser procedure. If a contact lens is used, your physician will instill an anaesthetic drop and place a small contact lens on your eye. Then you will place your head on the frame of the laser machine to keep your eye still, your physician will then focus the laser on the capsule.

The laser then makes small holes in the capsule, often in a circle. The holes join up and, like tearing a postage stamp, a small hole is made in the capsule. The segment drops to the bottom of the eye.

